

UGBforum



Literaturliste zum Beitrag:

Bewegungstherapie in der Onkologie: Sport tut gut

Dr. Thorsten Schmidt, UGBforum 4/23, S. 176-178

Adams SC, Segal RJ, McKenzie DC et al. (2016) Impact of resistance and aerobic exercise on sarcopenia and dynapenia in breast cancer patients receiving adjuvant chemotherapy. A multicenter randomized controlled trial. Breast Cancer Res Treat 158: 497–507.

Adamsen L, Quist M, Andersen C et al. (2009) Effect of a multimodal high intensity exercise intervention in cancer patients undergoing chemotherapy. Randomised controlled trial. BMJ 339: b3410.

Barnes B (2016) Bericht zum Krebsgeschehen in Deutschland 2016. Robert Koch-Institut, Berlin.

Baumann FT, Bloch W, Jäger E (2012) Sport und körperliche Aktivität in der Onkologie. Springer.

Buffart LM, Kalter J, Sweegers MG et al. (2017) Effects and moderators of exercise on quality of life and physical function in patients with cancer. An individual patient data meta-analysis of 34 RCTs. Cancer Treat Rev 52: 91–104.

Campbell KL, Winters-Stone KM, Wiskemann J et al. (2019) Exercise Guidelines for Cancer Survivors. Consensus Statement from International Multidisciplinary Roundtable. Med Sci Sports Exerc 51: 2375–2390.

Carayol M, Bernard P, Boiché J et al. (2013) Psychological effect of exercise in women with breast cancer receiving adjuvant therapy. What is the optimal dose needed? Ann Oncol 24: 291–300.

Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF (2017) Leitlinienprogramm Onkologie. S3-Leitlinie Früherkennung, Diagnose, Therapie und Nachsorge des Mammakarzinoms.

UGBforum



Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF (2018) S3-Leitlinie zur Diagnostik, Therapie und Nachsorge für Patienten mit einer chronischen lymphatischen Leukämie (CLL).

Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF (2021) Leitlinienprogramm Onkologie S-3 Leitlinie Prostatakarzinom.

Friedenreich CM, Shaw E, Neilson HK et al. (2017) Epidemiology and biology of physical activity and cancer recurrence. *J Mol Med* 95: 1029–1041.

Furmaniak AC, Menig M, Markes MH (2016) Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev* 9: CD005001.

Ganz PA, Kwan L, Castellon SA et al. (2013) Cognitive complaints after breast cancer treatments. Examining the relationship with neuropsychological test performance. *J Natl Cancer Inst* 105: 791–801.

Haydon AMM, Macinnis RJ, English DR et al. (2006) Effect of physical activity and body size on survival after diagnosis with colorectal cancer. *Gut* 55: 62–67.

Höh J-C, Schmidt T, Hübner J (2018) Physical activity among cancer survivors—what is their perception and experience? *Support Care Cancer* 26: 1471–1478.

Holmes MD, Chen WY, Feskanich D et al. (2005) Physical activity and survival after breast cancer diagnosis: 2479–2486.

Knols R, Aaronson NK, Uebelhart D et al. (2005) Physical exercise in cancer patients during and after medical treatment. A systematic review of randomized and controlled clinical trials. *J Clin Oncol* 23: 3830–3842.

Kuehr L, Wiskemann J, Abel U et al. (2014) Exercise in patients with non-small cell lung cancer. *Med Sci Sports Exerc* 46: 656–663.

Lahart IM, Metsios GS, Nevill AM et al. (2018) Physical activity for women with breast cancer after adjuvant therapy. *Cochrane Database Syst Rev* 1: CD011292.

UGBforum



Mishra SI, Scherer RW, Snyder C et al. (2012) Exercise interventions on health-related quality of life for people with cancer during active treatment. Cochrane Database Syst Rev: CD008465.

Quasthoff S, Hartung HP (2002) Chemotherapy-induced peripheral neuropathy. J Neurol 249: 9–17.

Schmidt T, Berner J, Jonat W et al. (2017) Influence of arm crank ergometry on development of lymphedema in breast cancer patients after axillary dissection. A randomized controlled trial. J Rehabil Med 49: 78–83.

Schmidt T, Weisser B, Dürkop J et al. (2015) Comparing Endurance and Resistance Training with Standard Care during Chemotherapy for Patients with Primary Breast Cancer. Anticancer Res 35: 5623–5629.

Schmitz KH, Ahmed RL, Troxel A et al. (2009) Weight lifting in women with breast-cancer-related lymphedema. N Engl J Med 361: 664–673.

Steindorf K, Schmidt ME, Klassen O et al. (2014) Randomized, controlled trial of resistance training in breast cancer patients receiving adjuvant radiotherapy. Results on cancer-related fatigue and quality of life. Ann Oncol 25: 2237–2243.

Streckmann F, Kneis S, Leifert JA et al. (2014) Exercise program improves therapy-related side-effects and quality of life in lymphoma patients undergoing therapy. Ann Oncol 25: 493–499.

Streckmann F, Lehmann HC, Balke M et al. (2018) Sensorimotor training and whole-body vibration training have the potential to reduce motor and sensory symptoms of chemotherapy-induced peripheral neuropathy-a randomized controlled pilot trial. Support Care Cancer.

van Waart H, Stuiver MM, van Harten WH et al. (2015) Effect of Low-Intensity Physical Activity and Moderate- to High-Intensity Physical Exercise During Adjuvant Chemotherapy on Physical Fitness, Fatigue, and Chemotherapy Completion Rates. Results of the PACES Randomized Clinical Trial. J Clin Oncol 33: 1918–1927.

UGBforum



Vollmers PL, Mundhenke C, Maass N et al. (2018) Evaluation of the effects of sensorimotor exercise on physical and psychological parameters in breast cancer patients undergoing neurotoxic chemotherapy. *J Cancer Res Clin Oncol* 144: 1785–1792.

Wiskemann J, Dreger P, Schwerdtfeger R et al. (2011) Effects of a partly self-administered exercise program before, during, and after allogeneic stem cell transplantation. *Blood* 117: 2604–2613.1. Friedrichsen H. Mechanismen bei oxidativem Stress und Entzündung. *Zeitschrift für Orthomolekulare Medizin* 2018; 16: 4–11