

Literaturliste zum Beitrag:

ADHS und eine oligoantogene Ernährung

Clement C, Fleischhaker C. ADHS und Ernährung. 6/16, S. 296

Bergmann K. (2014) ADHS und Ernährung: Mythen, Fakten, Zusammenhänge. Disserta Verlag, Hamburg.

Conners CK. (1997) Conners' Rating Scales-Revised technical manual. Multi-Health Systems; North Tonawanda, NY.

DuPaul GJ. (1991) Parent and teacher ratings of ADHD symptoms: psychometric properties in a community-based sample. J Clin Child Psychol 1991; 20: 242–253.

Egger J, Carter CM, Graham PJ, et al. (1985) Controlled trial of oligoantigenic treatment in the hyperkinetic syndrome. Lancet. 1(8428): 540-545.

Hiedl S. (2004) Duodenale VIP-Rezeptoren in der Dünndarmmukosa bei Kindern mit nahrungsmittelinduziertem hyperkinetischen Syndrom, Dissertation, LMU München.

Kaplan BJ, McNicol J, Conte RA, Moghadam HK. (1989) Dietary replacement in preschool-aged hyperactive boys. Pediatrics 83:7–17.

McCann D, Barrett A, Cooper A, et al. (2007) Food additives and hyperactive behaviour in 3-year-old and 8/9-year-old children in the community: a randomised, double-blinded, placebo-controlled trial. Lancet, 370:1560-1567.

Müller C. (2008) Verhaltensveränderungen unter einer standardisierten Eliminationsdiät bei Kindern mit Aufmerksamkeits-Defizit-Syndrom (ADS /ADHS): eine explorative Studie: Neue Akzente, 78: 8-12.

Pelsser LM, Frankena K, Toorman J, et al. (2009) A randomised controlled trial into the effects of food on ADHD. Eur Child Adolesc Psychiatry 18(1):12-19.

Pelsser LM, Frankena K, Toorman J, et al. (2011) Effects of a restricted elimination diet on the behaviour of children with attention-deficit hyperactivity disorder (INCA study): a randomised controlled trial. Lancet 377(9764): 494-503.

Pelsser LM, Buitelaar JK. (2002) Favourable effect of a standard elimination diet on the behavior of young children with attention deficit hyperactivity disorder (ADHD): a pilot study]. Ned Tijdschr Geneesk. 146(52):2543-2547.

Schab DW, Trinh NH. (2004) Do artificial food colors promote hyperactivity in children with hyperactive syndromes? A meta-analysis of double-blind placebo-controlled trials. J Dev Behav Pediatr 19(6):423-434.

Schulte-Körne G, Deimel W, Gutenbrunner C, et al. (1996) Effect of an oligo-antigen diet on the behavior of hyperkinetic children. Zeitschrift für Kinder- Jugendpsychiatrie und -psychotherapie 24(3): 176-183.

Shannon WR. (1922) Neuropathologic manifestations in infants and children as a result of anaphylactic reaction to foods contained in their dietary. American Journal of Disabled Children, 24, 89-94.

Sonuga-Barke EJ, Brandeis D, Cortese S, et al. (2013) Nonpharmacological interventions for ADHD: systematic review and meta-analyses of randomized controlled trials of dietary and psychological treatments. Am J Psychiatry 170(3):275-289.

Tarver J, Daley D, Sayal K. (2014) Attention-deficit hyperactivity disorder (ADHD): an updated review of the essential facts. Child Care Health Dev. 40(6):762-774.

Thapar A, Cooper M. (2016) Attention deficit hyperactivity disorder. Lancet. 387 (10024):1240-1250.

