

Literaturliste zum Beitrag:

Diabetes mellitus und seine Spätfolgen

Dr. Fischer B, UGB-Forum 2/12, S. 85

ALTIITUDE- Studie, www.akdae.de/Arzneimittelsicherheit/RHB/20120105.pdf

Boulton AJM, Vileikyte L, Ragnarson-Tennvall G, et al. (2005). The global burden of diabetic foot disease. Lancet 2005;366:1719-1724

H. Hauner (2012): Diabetesepidemie und Dunkelziffer. S. 8- 13 in Deutscher Gesundheitsbericht Diabetes 2012

Oh K, Hu FB, Manson JE, Stampfer MJ, Willett WC (2005) Dietary Fat Intake and Risk of Coronary Heart Disease in Women: 20 Years of Follow-up of the Nurses' Health Study. In: Am J Epidemiol. 2005 Apr 1;161(7):672-9.

Ohkubo Y, Kishikawa H, Araki E et al (1995). Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized controlled prospective 6-year study. Diabetes Res Clinical Pract 1995; 28: 103–17

Patterson CC, Dahlqvist GG, Gyüreüs E, Green A, Soltesz G for the EURODIAB Study Group (2009) Incidence trends for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–2020: a multicentre prospective registration study. Lancet, 2009, 373: 2027–2033

Praxisempfehlungen der Deutschen Diabetes-Gesellschaft (2011) Hasslacher C et al. Diabetische Nephropathie, Diabetologie 2011; 6: S111–S114

Qualitätssicherungsbericht 2006 (2008). Disease-Management-Programme in Nordrhein, Düsseldorf, 2008

The ADVANCE Collaborative Group (2008). Intensive Blood Glucose Control and Vascular Outcomes in Patients with Type 2 Diabetes. N Eng J Med (2008) 358:2560-2572

The Action to Control Cardiovascular Risk in Diabetes Study Group (2008). Effect of Intensive Glucose Lowering in Type 2 Diabetes. N Eng J Med (2008) 358:2545-2559

The Diabetes Control and Complications Trial Research Group (1993). The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. N Engl J Med 329:977–986, 1993

UK Prospective Diabetes Study (UKPDS) Group (1998). Intensive blood-glucose control with sulfonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). Lancet 1998; 352: 837–53

Whitmer et al (2009). Hypoglycemic Episodes and Risk of Dementia in older Patients with Typ 2 Diabetes, JAMA 2009, 301 (15): 1565-1572