

FODMAP-Diät bei Reizdarm erfolgreich

Beatrice Schilling, UGBforum 4/2018, Beatrice Schilling

Literaturliste zum Artikel

P. Wittkamp et al: Prävalenz des Reizdarmsyndroms nach den Rom-III-Kriterien in Deutschland und Zusammenhänge mit potentiellen Risikofaktoren, Zeitschrift für Gastroenterologie September 2012

Ziegler JU et al.: Wheat and the irritable bowel syndrome – FODMAP levels of modern and ancient species and their retention during bread making. Journal of Functional Foods 25 (2016), 257–266.

Biesiekierski JR et al.: No effects of gluten in patients with self-reported nonceliac gluten sensitivity after dietary reduction of fermentable, poorly absorbed, short-chain carbohydrates. Gastroenterology. 2013 Aug;145(2):320-8.e1-3.

Ong DK et al.: Manipulation of dietary short chain carbohydrates alters the pattern of gas production and genesis of symptoms in irritable bowel syndrome. J Gastroenterol Hepatol. 2010 Aug;25(8):1366-73.

Halmos EP et al.: A diet low in FODMAPs reduces symptoms of irritable bowel syndrome. Gastroenterology. 2014 Jan;146(1):67-75.e5.

Altobelli et al; Nutrients 2017,9,940;doi:10.3390 Low-FODMAP-diet improves Irritable Bowel Syndrom: A Meta-Analysis

Varju et al; PloS ONE 12(8)e0182942: Low FODMAP diet improves symptoms of adults suffering from IBS compared to standard diet: A Metaanalysis of clinical studies

Schumann D et al. Nutrition 2018 Jan; 45:24-31. doi: 10.1016/j.nut.2017.07.004. Low fermentable, oligo-, di, mono-saccharides and polyol diet in the treatment of irritable bowel syndrome: A systematic review and meta-analysis.