

Literaturliste zum Beitrag:

**Fettleber: ein unterschätztes Problem**

Prof. Dr. Werner O. Richter, UGBforum 4/15, S. 168

Bjermo HD et al (2012). Effects of n-6 PUFAs compared with SFAs on liver fat, lipoproteins, and inflammation in abdominal obesity: a randomized controlled trial. Am. J. Clin. Nutr. 95, 1003-1012

Lazo M et al (2010). Effect of a 12-month intensive lifestyle intervention on hepatic steatosis in adults with type 2 diabetes. Diabetes Care 33, 2156-2163

Petersson HJ et al (2010). Serum fatty acid composition and insulin resistance are independently associated with liver fat markers in elderly men. Diabetes Res. Clin. Pract. 87, 379-384

Visser ME et al (2011). Hepatic steatosis does not cause insulin resistance in people with familial hypobetalipoproteinemia. Diabetologia 54, 2113-2121

Westerbacke JK et al (2005). Dietary fat content modifies liver fat in overweight nondiabetic subjects. J. Clin. Endocrinol. Metab. 90, 2804-2809

Yki-Järvinen H (2010). Liver fat in the pathogenesis of insulin resistance and type 2 diabetes. Dig. Dis. 28, 203-209

Yki-Järvinen H (2010). Nutritional modulation of non-alcoholic disease and insulin resistance: human data. Curr. Opin. Clin. Nutr. Metab. Care 13, 709-714